

The Health-Boosting Properties of Super Foods

By Gary Null, PhD, and Martin Feldman, MD

In recent years, nutritional supplements have become increasingly high tech, providing physicians and their patients with advanced formulations for many healthcare needs. But despite their many benefits, these products should not detract from the more fundamental route to combating chronic disorders and improving one's health: the consumption of highly nutritious, powerhouse foods.

Like all healthy foods, “super foods” enhance a variety of bodily processes—but simply do it better. These foods contain high levels of antioxidants, phytochemicals, phytosterols, and dietary fiber that give them both preventive and therapeutic health properties. Native traditions throughout the world have long held that certain vegetables, fruits, and grains are especially powerful purveyors of health benefits. But it was not until these natural products were studied through modern biochemistry, botanical science, molecular biology, and clinical research that their extraordinary properties became more widely known.

In this article, we present 18 super foods that should be featured in the diet of healthcare providers and their patients. The benefits described have been culled from the

medical literature, and a sampling of the research conducted on these foods and their nutritional components is included in the references. What follows is a look at the specific preventive and curative properties of these super foods:

Apples

For thousands of years, apples (*malus sylvestris*) have been used to address numerous medical conditions, including diabetes, fevers, inflammatory disorders, and heart ailments. In addition to confirming many of the healthful properties of apples, modern research has identified invaluable phytochemicals contained in the fruits. One phytochemical found in apples is phloretin, a natural antibiotic. Apples also contain pectin and pectic acids that add essential bulk to a diet.

The apple's tannins, quercetin, alpha-farnesene, shikimic acid, and chlorogenic acid offer health benefits as well. By increasing the production of the neurotransmitter acetylcholine, for example, they help offset cognitive decline due to oxidative damage. Apples also have high levels of phenols and polyphenols and possess other antioxidant, chemoprotective properties. Consequently, they help guard against a variety of cancers, including leukemia and cancer of the colon, lung, breast, liver, and skin. These chemicals also provide essential nutrients that improve cardiovascular health, reduce the risk of coronary heart disease and stroke, and prevent atherosclerosis.

Apricots

This fruit had a long and rich history in the medical practices of China and India. In traditional Chinese medicine, apricots and their kernels are prescribed for the treatment of

asthma, cough, and constipation. The fruit is a stronghold of vitamin C, vitamin K, beta-carotene, thiamine, niacin, and iron. Japanese scientists have studied the ability of apricots to inhibit the pathogenic bacteria associated with ulcers and acute gastritis.

Bananas

Although low in calories, bananas provide essential nutrients such as vitamin B6, vitamin C, potassium, and manganese. They also stimulate probiotic activity, which sustains a healthy gut flora. Bacteria in the gastrointestinal system are critical for the proper digestion and absorption of nutrients. Bananas help keep this system on track. Recent findings have indicated that bananas may offer protection against kidney cancer, particularly in women, and aid renal function.

Blueberries

Many berries have health-boosting properties. The black, blue, and red varieties are especially known for the antioxidants they contain. Blueberries, in particular, have flavonoids, phenolic and polyphenol compounds, all of which have shown some ability to reverse cellular aging of cognitive and motor functions. In one recent study of the antioxidant levels of 100 foods, blueberries scored highest.

Other studies have shown that blueberries protect brain health, improve memory, and sustain coordination by, for one, enhancing communication between nerve cells. This activity provided protection against serious neurodegenerative diseases, such as dementia and Alzheimer's. Blueberries also have anti-inflammatory properties that protect the skin, the joints, and the cardiovascular and neurological systems. The consumption of

blueberries has proven beneficial to people with diabetes. This fruit also prevents bone loss and inhibits cancer cell proliferation, particularly in prostate and colon cancer.

Broccoli

Broccoli is a super food because of its high concentration of two phytochemicals—diindolymethane and isothiocyanate—that are powerful immunomodulators. Broccoli strengthens the immune system, which means it helps in the fight against cancer (particularly breast and prostate cancer) and boosts the body's protection against bacterial and viral infections. Broccoli contains other anticancer agents as well, such as glucoraphanin. Due to these observed properties, a substantial amount of research is being conducted on broccoli's mutagenic qualities.

This vegetable is rich in vitamins A, B5, B6, B9 (folate), C, and K and in dietary fiber. It provides moderate amounts of calcium, iron, phosphorus, and potassium. As with other leafy green vegetables, broccoli contains lutein and zeaxanthin, which foster eye health. Because it has more calcium than even most dairy products, broccoli can protect bones and increase bone mass.

Carrots

Carrots are a chief provider of carotenoids, a family of antioxidants proven to block DNA and cellular membrane damage caused by free radicals. Carrots are rich in alpha-carotene and lycopene, phytochemicals that have anti-carcinogenic properties, especially in relation to colon, lung, prostate, and stomach cancers. The lesser-known black and purple

carrots have high levels of anthocyanin, a powerful anticancer biochemical that has been found to slow cancer cell proliferation by as much as 80 percent.

The long-held belief that carrots improve vision is supported by their high content of retinoids that benefit ocular health. Carrots also have been shown to boost brain function and provide cardiovascular benefits, such as decreasing cholesterol. Diabetics should keep carrots in their diet because they are a good source of vitamin A, which lowers blood sugar and aids in the development of insulin-producing cells in the pancreas. One cup of raw carrots can provide nearly 700 percent of the recommended daily intake of vitamin A and 220 percent of vitamin K, which is critical for bone health.

Garlic

While garlic contains phytonutrients similar to those found in onions, it also possesses selenium, a substance that, according to some studies, offers protection against various cancers and the deterioration of the body caused by free radicals. Researchers have studied the ability of garlic to guard against heart disease and arterial calcification (hardening of the arteries) and to reduce cholesterol and blood pressure. Because it is a source of the flavonoid quercetin, garlic contains antibiotic properties that empower it to fight colds, stomach viruses, and yeast infections.

Ginger

Ginger is used throughout the world to cure dyspepsia (stomach upsets), reduce gastrointestinal gases, and relieve nausea caused by pregnancy, seasickness, and even drugs used in chemotherapy. Ginger is composed largely of fragrant essential oils that

give it a distinctive aromatic flavor. One of these oils, gingerol, makes it a natural sedative for calming the gastrointestinal tract. This oil also provides some protection against pathogenic bacteria that upset the stomach. Ginger is rich in antibiotic properties that combat the GI infections which cause diarrhea and dehydration.

Folk medicine has long honored ginger. While some scientists may dismiss folk medicine, it should be remembered that many modern pharmaceuticals were derived from folk remedies and then price tagged. This folk science, now supported by modern science, has viewed ginger as a mild immune booster that wards off colds, flus, sinus congestion, and coughs. New evidence suggests that ginger helps to lower cholesterol. Preliminary findings in animal studies also suggest that ginger may help to treat diabetes.

Goji Berry

Also known as wolfberry in its native Europe, the goji plant is found in much of Asia, where it appears in exotic (to Westerners) Tibetan and Himalayan descriptions. The word *goji* is actually a Westernization of the Chinese word for the berry, which can be transliterated as “gouqi.” The berry is a common ingredient in traditional Chinese medicine, dating back thousands of years in use.

The oblong red goji berry easily fulfills the requirements of a super food. It has a high concentration of phytochemicals, amino acids, vitamins B and C, and beta-carotene. Additionally, it contains 11 essential and 22 trace dietary minerals, is an outstanding source of the antioxidant lycopene, and is moderately high in alpha-linolenic acid. The goji berry also can supply extra protein, dietary fiber, calcium, zinc, and selenium.

This nutritional profile gives the goji berry many health-enhancing properties. This fruit protects against cardiovascular and inflammatory diseases and age-related vision disorders (such as glaucoma and macular degeneration). Studies have pointed to the berry's neuro-protective, immunomodulatory, and anticancer properties as well. This last benefit was underscored by a study published in the Chinese Journal of Oncology, which indicated that cancer patients responded better to treatment while on a diet that included goji. However, the study recommended that individuals on blood-thinning medications avoid eating goji berries, which may interfere with the drugs. Finally, the goji berry offers liver protection and can improve sexual function.

Green Tea

The ingredient in tea—and particularly green tea—that has stirred the most scientific interest is catechin. Approximately 25 percent of a dry tea leaf is catechin. Although traces of catechin are also found in chocolate, wine, and other fruits and vegetables, it is tea that offers the greatest amount of this super nutrient.

The multi-tasking catechin has been shown to reduce the plaque buildup of atherosclerosis, protect against infectious bacteria, and reduce oxidative stress. Tea catechins are especially important in a polluted world because they can improve DNA replication and protect against genetic damage from environmental toxins. Recent studies have noted the anti-inflammatory properties of catechin and suggested it can play a role in battling cancer. Other research has noted that green tea can improve bone density and cognitive function, reduce the risk of developing kidney stones, and strengthen heart

function. There is some evidence that the polyphenols of green tea protect against the brain cell death associated with Parkinson's and Alzheimer's diseases.

Legumes

Individuals who consume a Western diet, especially in America, ignore the nutritional value of most legumes to their own detriment. This category of super food includes not only beans, peas and lentils—the foods most commonly identified as legumes—but also alfalfa, clover, peanuts, and cashews.

These vegetables and grains are excellent sources of dietary fiber, which reduces cholesterol and helps manage blood sugar levels. One cup of lentils can provide upwards of 65 percent of the minimum daily requirement for fiber. Given this high fiber content, the frequent consumption of legumes will enhance gastrointestinal and colon health.

Legumes contain energy-boosting protein and iron. Looking at specific entities in this group, black beans are rich in the potent antioxidant anthocyanidins, which promotes heart and vascular health. Green beans are excellent sources of vitamins C and K. Garbanzo beans, commonly known as chickpeas, are a superb source of molybdenum, which strengthens teeth and preserves tooth enamel.

Another important legume that is not well-known in the U.S. is the adzuki bean. Originally from the Himalayas and standard in East Asian cooking, adzuki beans are a rich source of magnesium, potassium, iron, zinc, and B vitamins. Very high in soluble fiber, the adzuki helps eliminate bad cholesterol from the body. In Japan, it is treasured for its kidney and bladder health-promoting function and is used in weight-loss programs.

Combining legumes with whole grains will maximize their benefits. Legumes are high in lysine but very low in methionine, an essential amino acid that supports cellular life, while whole grains are replete with this amino acid but low in lysine. A wholesome, integrated vegetarian diet will contain a balance of legumes and grains.

Leafy Vegetables

This group of super foods includes spinach, kale, arugula, Swiss chard, cabbage, collard greens, and watercress. One feature common to the dark green leafy vegetables is that they are high in carotenoids and other antioxidants that guard against heart disease, cancer, and problems with blood sugar regulation. Beyond that, each leafy green offers its own health benefits, so a healthy diet should contain all members of this category.

For example, one cup of cooked kale provides over 1,300 percent of the daily requirement of vitamin K needed for maximum bone health. It is also rich in calcium and manganese, other nurturers of bone density. Like broccoli, kale also contains the anticancer phytochemical sulforaphane.

Cabbage contains glutamine, an amino acid that contributes to anti-inflammatory activities. It also protects against infectious complications due to human papillomavirus (HPV). The juice from cabbage will speed up the healing of peptic ulcers.

Spinach is one good source of dietary iron. Per gram, it generally contains over 30 percent more iron than a hamburger does. (Any diet heavy in spinach should include sufficient vitamin C to help assimilate the iron.) Spinach also ia an excellent source of folic acid, calcium, copper, zinc, and selenium.

Watercress is a superb source of phytochemicals. It serves as a diuretic and digestive aid, helps protect against lung cancer, and strengthens the thyroid. Collard greens supply ample quantities of the immune response modulator diindolylmethane.

Mushrooms

A wealth of peer-reviewed literature shows that many edible mushrooms are among the more important immune builders in the plant kingdom. In particular, medicinal mushrooms inhibit tumor growth, strengthen immunity, and have anti-pathogenic and blood-sugar-lowering properties.

Among approximately 200 varieties of mushrooms whose health-enhancing skills have been noted are the chaga, cordyceps, maitake, oyster, ortobello, reishi, shiitake, and turkey tail. Although all of these types can be obtained in fresh or dried form, shiitake mushrooms currently are the easiest to obtain in the United States.

A list of the health benefits of mushrooms would have to include their antiviral and antibacterial properties, which in different varieties have shown some effectiveness against pathogens including polio, hepatitis B, influenza, candida, Epstein-Barr virus, streptococcus, and tuberculosis. The scientific literature also discusses the mutagenic benefits of mushrooms, which can be enlisted in the fight against leukemia, sarcoma, and the bladder, breast, colon, liver, lung, prostate, stomach cancers, even in advanced stages.

Onions

This super food is rich in vitamin B6, vitamin C, manganese, molybdenum (essential in preserving tooth enamel), potassium, phosphorous, and copper. Onions are an excellent

source of quercetin, which works with vitamin C to help the body eliminate bacteria and strengthen immunity. A rule of thumb is that the more pungent the onion, the greater its health benefits.

Onions are particularly important in the diet of diabetics because they are rich in chromium, a trace mineral that helps cells respond to insulin. Moreover, refined sugar depletes the body's chromium levels, so onions are an excellent source of chromium replacement for anyone who consumes refined sugar.

Onions also help to reduce blood pressure and cholesterol and strengthen bone health. They have anti-inflammatory benefits, reducing symptoms related to inflammatory conditions such as asthma, arthritis, and respiratory congestion. Some studies have noted that onions lessen the adverse effects of colds and flus.

Oranges

The orange is a vitamin- and mineral-packed fruit, rich in vitamins A, B, and C and potassium and calcium. It is an excellent source of fiber as well. One phytonutrient in oranges that places it in the super food category is the flavonoid hesperidin. This biochemical helps support healthy blood vessels and reduces cholesterol.

The orange's defining health trait is its high content of vitamin C, an important antioxidant that limits free radicals while also building the immune system. Vitamin C's healing properties are well known and have been repeatedly scientifically validated. These include lessening arterial plaque and protecting against Alzheimer's, Parkinson's, and Crohn's diseases, arthritis, and diabetes.

Peppers (capsicum)

Native American folk medicine, which has so many features we can learn from, gave a prominent place in its pharmacology to peppers of the capsicum family (including bell and chili peppers). Recent work suggests that the nutrient capsaicin, contained in these peppers, is a natural analgesic and a neuro-inflammatory blocker that relieves aches and pains in joints and muscles. This is one reason why Native American medicine prescribed a topical application of pepper to painful areas of the body.

Adding to the super food designation of peppers is promising research in Canada that has explored the uses of capsaicin in the treatment of type 1 diabetes. Other research has found benefits for individuals with prostate cancer and leukemia. Some scientists have noted that the much-studied capsaicin helps with weight loss, stimulation of insulin-producing cells, and prevention of LDL cholesterol oxidation. Another benefit recently uncovered is that capsaicin protects against stomach ulcerations and induces apoptosis (cancer cell death) in lung cancer.

In addition to capsaicin, peppers are rich in the antioxidant vitamin A, vitamins B1, B6, E, and K, and potassium, magnesium, and iron. Yellow peppers are rich in lutein and zeaxanthin, which protect against eye disease and blindness.

Tomatoes

All of the super foods contain highly potent organic compounds, such as phytochemicals, that boost their health-giving properties. Tomatoes are no exception. They are the best source of lycopene, a carotenoid biochemical that gives tomatoes their red color and is

packed with healthful properties. An estimated 80 percent of the lycopene consumed in the United States is derived from tomatoes and tomato-based foods.

There is a vast body of scientific literature confirming lycopene's antioxidant and anti-mutagenic properties. This chemical is noteworthy for its protection against and treatment of various cancers, including those of the bladder, breast, cervix, lungs, mouth, ovary, prostate, and stomach. Because diabetics often have low levels of lycopene in their blood, tomatoes should be a regular part of their diets.

Tomatoes have been shown to prevent cholesterol oxidation, lower blood pressure, and decrease the risk of atherosclerosis. Other benefits that may accrue to the eater of these plants is improved renal function. Tomatoes have anti-viral and anti-bacterial qualities. In particular, lycopene can protect against human papillomavirus, a pathogen that has been associated with cancer.

Tomatoes are rich in most of the B complex vitamins, potassium, manganese, chromium, folate, and iron. They also are an excellent source of the amino acid tryptophan, which is important for neurological health and can improve sleep.

Whole Grains

By now most Americans are aware that whole grain breads and pastas are healthier than those made from white flour, and brown rice is higher in nutrients and health benefits than is white rice. However, once a person has changed over to brown rice and whole grain breads, he or she still has a rich world of whole grains to explore, each of which offers unique health benefits and phytonutrients.

Like legumes, whole grains are rich in fiber. One grain, spelt, is used in breads and pastas and provides 75 percent of the recommended daily requirement for vitamin B2. Spelt is highly water soluble, which means its nutrients are easily absorbed. There is evidence that spelt is a good choice for diabetics. Another grain, barley, is distinguished by being an excellent source of selenium, a substance that reduces the risk of colon disorders and colorectal cancer. Because barley is high in tryptophan, it will aid in sleep regulation. A third important grain, millet, is high in manganese, magnesium, and phosphorous, all of which support cardiovascular health.

Two less familiar grains are kamut and quinoa. The Glycemic Research Institute in Washington, D.C., has trumpeted kamut for its low-glycemic properties, which makes it an ideal super food for diabetics, athletes, and people suffering from obesity. It also is an excellent substitute for those with wheat allergies because it has 65 percent more amino acids than wheat.

Quinoa has been identified as a super food among grains because of its ability to balance blood sugar and provide high-quality fiber and protein in the diet. It is higher in calcium, phosphorus, iron, and zinc than are wheat, barley and corn. Quinoa is one of the most complete foods in nature, earning its super-food status not only for the properties described above but also its role in protecting against atherosclerosis and breast cancer and, as a probiotic, fostering beneficial micro-flora in the gut.

Correspondence:

Gary Null, PhD
2307 Broadway
New York, New York 10024 USA

Martin Feldman, MD
132 East 76th Street
New York, New York 10021 USA
e-mail: precisemd@aol.com

The Authors

Gary Null, PhD, has authored more than 60 books on health and nutrition and numerous articles published in research journals. He is Adjunct Professor, Graduate Studies, Public Health Curriculum, at Fairleigh Dickinson University in Teaneck, N.J. Null holds a Ph.D. in human nutrition and public health science from the Union Graduate School.

Martin Feldman, MD, practices complementary medicine. He is an Assistant Clinical Professor of Neurology at the Mount Sinai School of Medicine in New York City.

REFERENCES

Apples

Boyer J, Liu RH. Apple phytochemicals and their health benefits *Nutr J* 2004; 3:5.
 Jung M, Triebel S, Anke T, Richling E, Erkel G. Influence of apple polyphenols on inflammatory gene expression. *Mol Nutr Food Res* 2009; 53(10):1263-80.

Wojdyło A, Oszmiański J, Laskowski P. Polyphenolic compounds and antioxidant activity of new and old apple varieties. *J Agric Food Chem* 2008; 56(15):6520-30.

Apricots

Enomoto S, Yanaoka K, Utsunomiya H, et al. Inhibitory effects of Japanese apricot (*Prunus mume* Siebold et Zucc.; Ume) on *Helicobacter pylori*-related chronic gastritis. *Eur J Clin Nutr* 2010; 64(7):714-9.

Fujita K, Hasegawa M, Fujita M, et al. [Anti-*Helicobacter pylori* effects of Bainiku-ekisu (concentrate of Japanese apricot juice)] *Nippon Shokakibyō Gakkai Zasshi* 2002; 99(4):379-85. [Article in Japanese]

Bananas

Chow J. Probiotics and prebiotics: A brief overview. *J Ren Nutr* 2002; 12(2):76-86.

Rashidkhani B, Lindblad P, Wolk A. Fruits, vegetables and risk of renal cell carcinoma: a prospective study of Swedish women. *Int J Cancer* 2005; 113(3):451-5.

Blueberries

Joseph JA, Shukitt-Hale B, Casadesus G. Reversing the deleterious effects of aging on neuronal communication and behavior: beneficial properties of fruit polyphenolic compounds. *Am J Clin Nutr* 2005; 81(1 Suppl):313S-316S.

Krikorian R, Shidler MD, Nash TA, et al. Blueberry supplementation improves memory in older adults. *J Agric Food Chem* 2010; 58(7):3996-4000.

Paul S, DeCastro AJ, Lee HJ, et al. Dietary intake of pterostilbene, a constituent of blueberries, inhibits the beta-catenin/p65 downstream signaling pathway and colon carcinogenesis in rats. *Carcinogenesis* 2010; 31(7):1272-8.

Torri E, Lemos M, Caliarì V, et al. Anti-inflammatory and antinociceptive properties of blueberry extract (*Vaccinium corymbosum*). *J Pharm Pharmacol* 2007; 59(4):591-6.

Broccoli

Kim HJ, Barajas B, Wang M, Nel AE. Nrf2 activation by sulforaphane restores the age-related decrease of T(H)1 immunity: role of dendritic cells. *J Allergy Clin Immunol* 2008; 121(5):1255-1261.e7.

Le HT, Schaldach CM, Firestone GL, Bjeldanes LF. Plant-derived 3,3'-Diindolylmethane is a strong androgen antagonist in human prostate cancer cells. *Biol Chem* 2003; 278(23):21136-45.

Li Y, Zhang T, Korkaya H, et al. Sulforaphane, a dietary component of broccoli/broccoli sprouts, inhibits breast cancer stem cells. *Clin Cancer Res* 2010; 16(9):2580-90.
Murugan SS, Balakrishnamurthy P, Mathew YJ. Antimutagenic effect of broccoli flower head by the ames salmonella reverse mutation assay. *Phytother Res* 2007; 21(6):545-7.

Carrots

Coleman AL, Stone KL, Kodjebacheva G, et al. Glaucoma risk and the consumption of fruits and vegetables among older women in the study of osteoporotic fractures. *Am J Ophthalmol* 2008; 145(6):1081-9.

Fernandes I, Faria A, Azevedo J, et al. Influence of anthocyanins, derivative pigments and other catechol and pyrogallol-type phenolics on breast cancer cell proliferation. *J Agric Food Chem* 2010; 58(6):3785-92.

Khachik F, Beecher GR, Smith JC Jr. Lutein, lycopene, and their oxidative metabolites in chemoprevention of cancer. *J Cell Biochem Suppl* 1995; 22:236-46.

Nicolle C, Cardinault N, Aprikian O, et al. Effect of carrot intake on cholesterol metabolism and on antioxidant status in cholesterol-fed rat. *Eur J Nutr* 2003; 42(5):254-61.

Nurk E, Refsum H, Drevon CA, et al. Cognitive performance among the elderly in relation to the intake of plant foods. The Hordaland Health Study. *Br J Nutr* 2010; 104(8):1190-201.

Garlic

Duda G, Suliburska J, Pupek-Musialik D. Effects of short-term garlic supplementation on lipid metabolism and antioxidant status in hypertensive adults. *Pharmacol Rep* 2008; 60(2):163-70.

Gorinstein S, Jastrzebski Z, Namiesnik J, et al. The atherosclerotic heart disease and protecting properties of garlic: contemporary data. *Mol Nutr Food Res* 2007; 51(11):1365-81.

Yeh YY, Liu L. Cholesterol-lowering effect of garlic extracts and organosulfur compounds: human and animal studies. *J Nutr* 2001; 131(3s):989S-93S.

Ginger

Ali BH, Blunden G, Tanira MO, Nemmar A. Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food Chem Toxicol* 2008; 46(2):409-20.

Alizadeh-Navaei R, Roozbeh F, Saravi M, et al. Investigation of the effect of ginger on the lipid levels. A double blind controlled clinical trial. *Saudi Med J* 2008; 29(9):1280-4.

Park M, Bae J, Lee DS. Antibacterial activity of [10]-gingerol and [12]-gingerol isolated from ginger rhizome against periodontal bacteria. *Phytother Res* 2008; 22(11):1446-9.

Priya Rani M, Padmakumari KP, Sankarikutty B, et al. Inhibitory potential of ginger extracts against enzymes linked to type 2 diabetes, inflammation and induced oxidative stress. *Int J Food Sci Nutr* 2010 Sep 28. [Epub ahead of print]

Goji Berry

Amagase H, Sun B, Borek C. *Lycium barbarum* (goji) juice improves in vivo antioxidant biomarkers in serum of healthy adults. *Nutr Res* 2009 Jan; 29(1):19-25.

Amagase H, Sun B, Nance DM. Immunomodulatory effects of a standardized *Lycium barbarum* fruit juice in Chinese older healthy human subjects. *J Med Food* 2009; 12(5):1159-65.

Cao GW, Yang WG, Du P. [Observation of the effects of LAK/IL-2 therapy combining with *Lycium barbarum* polysaccharides in the treatment of 75 cancer patients]. [Article in Chinese] *Zhonghua Zhong Liu Za Zhi* 1994;16(6):428-31.

Potterat O. Goji (*Lycium barbarum* and *L. chinense*): Phytochemistry, pharmacology and safety in the perspective of traditional uses and recent popularity. *Planta Med* 2010; 76(1):7-19.

Green Tea

Cooper R, Morr  DJ, Morr  DM. Medicinal benefits of green tea: Part I. Review of noncancer health benefits. *J Altern Complement Med* 2005; 11(3):521-8.

Dou QP. Molecular mechanisms of green tea polyphenols. *Nutr Cancer* 2009; 61(6):827-35.

Mandel SA, Avramovich-Tirosh Y, Reznichenko L, et al. Multifunctional activities of green tea catechins in neuroprotection. Modulation of cell survival genes, iron-dependent oxidative stress and PKC signaling pathway. *Neurosignals* 2005; 14(1-2):46-60.

Legumes

Itoh T, Furuichi Y. Lowering serum cholesterol level by feeding a 40% ethanol-eluted fraction from HP-20 resin treated with hot water extract of adzuki beans (*Vigna angularis*) to rats fed a high-fat cholesterol diet. *Nutrition* 2009; 25(3):318-21.

Kim JM, Kim JS, Yoo H, et al. Effects of black soybean [*Glycine max* (L.) Merr.] seed coats and its anthocyanidins on colonic inflammation and cell proliferation in vitro and in vivo. *J Agric Food Chem* 2008; 56(18):8427-33.

Yang Y, Zhou L, Gu Y, et al. Dietary chickpeas reverse visceral adiposity, dyslipidaemia and insulin resistance in rats induced by a chronic high-fat diet. *Br J Nutr* 2007; 98(4):720-6.

Leafy Vegetables

Carter P, Gray LJ, Troughton J, et al. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ* 2010; 341:c4229. doi: 10.1136/bmj.c4229.

Hecht SS, Chung FL, Richie JP Jr, et al. Effects of watercress consumption on metabolism of a tobacco-specific lung carcinogen in smokers. *Cancer Epidemiol Biomarkers Prev* 1995; 4(8):877-84.

Kim SY, Yoon S, Kwon SM, et al. Kale juice improves coronary artery disease risk factors in hypercholesterolemic men. *Biomed Environ Sci* 2008; 21(2):91-7.

Riby JE, Xue L, Chatterji U, et al. Activation and potentiation of interferon-gamma signaling by 3,3'-diindolylmethane in MCF-7 breast cancer cells. *Mol Pharmacol* 2006; 69(2):430-9.

Mushrooms

Akihisa T, Franzblau SG, Tokuda H, et al. Antitubercular activity and inhibitory effect on Epstein-Barr virus activation of sterols and polyisoprenepolyols from an edible mushroom, *Hypsizygus marmoreus*. *Biol Pharm Bull* 2005; 28(6):1117-9.

Akramiene D, Kondrotas A, Didziapetriene J, Kevelaitis E. Effects of beta-glucans on the immune system. *Medicina (Kaunas)* 2007; 43(8):597-606.

Gao L, Sun Y, Chen C, et al. Primary mechanism of apoptosis induction in a leukemia cell line by fraction FA-2-b-ss prepared from the mushroom *Agaricus blazei* Murill. *Braz J Med Biol Res* 2007; 40(11):1545-55.

Lu X, Chen H, Dong P, et al. Phytochemical characteristics and hypoglycaemic activity of fraction from mushroom *Inonotus obliquus*. *J Sci Food Agric* 2010; 90(2):276-80.

Wasser SP. Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides. *Appl Microbiol Biotechnol* 2002; 60(3):258-74.

Onions

Arai Y, Watanabe S, Kimura M, et al. Dietary intakes of flavonols, flavones and isoflavones by Japanese women and the inverse correlation between quercetin intake and plasma LDL cholesterol concentration. *J Nutr* 2000;130(9):2243-50.

Kook S, Kim GH, Choi K. The antidiabetic effect of onion and garlic in experimental diabetic rats: meta-analysis. *J Med Food* 2009;12(3):552-60.

Mayer B, Kalus U, Grigorov A, et al. Effects of an onion-olive oil maceration product containing essential ingredients of the Mediterranean diet on blood pressure and blood fluidity. *Arzneimittelforschung* 2001;51(2):104-11.

Wagner H, Dorsch W, Bayer T, et al. Antiasthmatic effects of onions: inhibition of 5-lipoxygenase and cyclooxygenase in vitro by thiosulfinates and "Cepaenes." *Prostaglandins Leukot Essent Fatty Acids* 1990; 39(1):59-62.

Oranges

Heo HJ, Choi SJ, Choi SG, et al. Effects of banana, orange, and apple on oxidative stress-induced neurotoxicity in PC12 cells. *J Food Sci* 2008; 73(2):H28-32.

Jin YR, Han XH, Zhang YH, et al. Antiplatelet activity of hesperetin, a bioflavonoid, is mainly mediated by inhibition of PLC-gamma2 phosphorylation and cyclooxygenase-1 activity. *Atherosclerosis* 2007; 194(1):144-52.

Kurowska EM, Spence JD, Jordan J, et al. HDL-cholesterol-raising effect of orange juice in subjects with hypercholesterolemia. *Am J Clin Nutr* 2000; 72(5):1095-100.

Peppers (Capsicum)

Ahuja KD, Ball MJ. Effects of daily ingestion of chilli on serum lipoprotein oxidation in adult men and women. *Br J Nutr*. 2006; 96(2):239-42.

Luo XJ, Peng J, Li YJ. Recent advances in the study on capsaicinoids and capsinoids. *Eur J Pharmacol* 2011; 650(1):1-7.

Mori A, Lehmann S, O'Kelly J, et al. Capsaicin, a component of red peppers, inhibits the growth of androgen-independent, p53 mutant prostate cancer cells. *Cancer Res*. 2006; 66(6):3222-9.

Schumacher MA. Transient receptor potential channels in pain and inflammation: therapeutic opportunities. *Pain Pract* 2010; 10(3):185-200.

Tomatoes

Bhuvanewari V, Nagini S. Lycopene: a review of its potential as an anticancer agent. *Curr Med Chem Anticancer Agents* 2005; 5(6):627-35.

Rodríguez-Muñoz E, Herrera-Ruiz G, Pedraza-Aboytes G, Loarca-Piña G. Antioxidant capacity and antimutagenic activity of natural oleoresin from greenhouse grown tomatoes (*Lycopersicon esculentum*). *Plant Foods Hum Nutr* 2009; 64(1):46-51.

Sedjo RL, Papenfuss MR, Craft NE, Giuliano AR. Effect of plasma micronutrients on clearance of oncogenic human papillomavirus (HPV) infection (United States). *Cancer Causes Control* 2003 May; 14(4):319-26.

Silaste ML, Alfthan G, Aro A, Kesäniemi YA, Hörkö S. Tomato juice decreases LDL cholesterol levels and increases LDL resistance to oxidation. *Br J Nutr* 2007 Dec; 98(6):1251-8.

Whole Grains

Murtaugh MA, Jacobs DR Jr, Jacob B, et al. Epidemiological support for the protection of whole grains against diabetes. *Proc Nutr Soc* 2003; 62(1):143-9.

Lee SH, Chung IM, Cha YS, Park Y. Millet consumption decreased serum concentration of triglyceride and C-reactive protein but not oxidative status in hyperlipidemic rats. *Nutr Res* 2010; 30(4):290-6.

Okarter N, Liu RH. Health benefits of whole grain phytochemicals. *Crit Rev Food Sci Nutr* 2010; 50(3):193-208.

Paško P, Zagrodzki P, Bartoń H, et al. Effect of quinoa seeds (*Chenopodium quinoa*) in diet on some biochemical parameters and essential elements in blood of high fructose-fed rats. *Plant Foods Hum Nutr* 2010; 65(4):333-8.

Slavin J. Why whole grains are protective: biological mechanisms. *Proc Nutr Soc* 2003; 62(1):129-34.